

String structures associated to indefinite Lie groups

Hisham Sati and Hyung-bo Shim

April 9, 2015

Abstract

String structures have played an important role in algebraic topology, via elliptic genera and elliptic cohomology, in differential geometry, via the study of higher geometric structures, and in physics, via partition functions. We extend the description of String structures from connected covers of the definite-signature orthogonal group $O(n)$ to the indefinite-signature orthogonal group $O(p, q)$, i.e. from the Riemannian to the pseudo-Riemannian setting. This requires that we work at the unstable level, which makes the discussion more subtle than the stable case. Similar, but much simpler, constructions hold for other noncompact Lie groups such as the unitary group $U(p, q)$ and the symplectic group $Sp(p, q)$. This extension provides a starting point for an abundance of constructions in (higher) geometry and applications in physics.

Contents

1	Introduction	2
2	The Postnikov tower, Whitehead tower, and variants	4
2.1	The Postnikov tower	4
2.2	The Whitehead tower	5
2.3	A variant view	7
2.4	Computational aspects	8
2.5	Special case: higher connected covers of classifying spaces and obstructions	10
3	Applications	13
3.1	Special orthogonal groups $SO(n)$ and $SO(p, q)$	13
3.2	Indefinite Spin groups	14
3.3	Indefinite String groups	17
3.4	String structure associated to indefinite unitary and symplectic groups	22
3.5	Relation to twisted structures	24

1 Introduction

Lie groups play an important role in characterizing symmetries. Picking the appropriate Lie group allows for certain structures to be unambiguously defined. For example, a Riemannian structure on a manifold requires the principal frame bundle corresponding to the tangent bundle to have an orthogonal group as a structure group. To talk about orientations one needs the special orthogonal group, and to properly discuss spinors one needs to lift to the double cover, which is the Spin group. All of these structures are low degree phenomena which can be encoded uniformly and succinctly via the Whitehead tower of the orthogonal group (see [27] [28] [26]).

In algebraic topology, one usually studies the Whitehead tower of the stable orthogonal group (see [27]). In particular, killing the third homotopy group leads to the stable String group (see [30] [31]). There are many constructions and applications associated with String structures and to the String group in various areas of mathematics and physics. The following is a sample which is necessarily incomplete. In algebraic topology, String structures play a role of orientation for elliptic cohomology [1] [30] [31]. In differential geometry, String connections play a role in geometrically characterizing bundles with the String group as a structure group [28] [23] [35] [6] [7]. In mathematical physics, conditions for having String structures arise as anomaly cancellation conditions [14] [27] [28].

This paper aims to initiate a new angle on the subject. We will be interested in the case of the indefinite orthogonal group $O(p, q)$, which from a geometric point of view can be viewed as the structure group of the tangent bundle of a pseudo-Riemannian manifold of dimension $n = p + q$. As in the Riemannian case, one is interested in considering oriented and then Spin pseudo-Riemannian manifolds and for that one needs to lift $O(p, q)$ to appropriate groups $SO(p, q)$ and $Spin(p, q)$, respectively. This involves a lot of subtleties and unlike the Riemannian case, here it is a priori not obvious which homotopy (su-)groups to kill in order to get to the appropriate covering group. We spend some time discussing this before embarking on considering the corresponding indefinite String groups. In geometry and physics, the cases $p = 1, 2$ are particularly interesting, as these correspond to the Lorentz group and the conformal group, respectively. We address all cases, and for the most part there is a natural split into cases $p = 1$, $p = 2$ and $p \geq 3$.

Another subtle matter in our discussion is that we need to work in the unstable range for the orthogonal groups, that is with $O(n)$ for finite n , i.e. without taking a limit on the rank as usually is done in the literature. For the Riemannian case, the Whitehead tower (cf. [27]) would be replaced by one involving finite n

$$\begin{array}{ccccc}
 & & BString(n) & & (1.1) \\
 & \nearrow & \downarrow & & \\
 & & BSpin(n) & \xrightarrow{H^4(BSpin(n); \mathbb{Z})} & K(\mathbb{Z}, 4) \\
 & \nearrow & \downarrow & & \\
 & & BSO(n) & \xrightarrow{H^2(BSO(n); \mathbb{Z}/2)} & K(\mathbb{Z}/2, 2) \\
 & \nearrow & \downarrow & & \\
 X & \xrightarrow{\quad} & BO(n) & \xrightarrow{H^1(BO(n); \mathbb{Z}/2)} & K(\mathbb{Z}/2, 1) .
 \end{array}$$

This gives a homotopy theoretical construction of 8-connected cover $(BO(n))\langle 8 \rangle$ of the classifying space of $O(n)$. Having a classifying space points to presence of a group. Indeed, Stolz [30] constructs a topological group $\widehat{G}$ as an extension

$$1 \rightarrow \text{Gauge}(P) \rightarrow \widehat{G} \rightarrow \text{Spin}(n) \rightarrow 1, \quad (1.2)$$

where $P \rightarrow \text{Spin}(n)$ is the $PU(\mathcal{H})$ -principal bundle with $\mathcal{H}$ an infinite dimensional separable Hilbert space such that $B\widehat{G}$ is the 8-connected cover of $BO(n)$. This topological group $\widehat{G}$ is homotopy equivalent to $\text{String}(n)$ so we can say $\text{String}(n)$ has a group structure. This also has a differentiable structure. Since then many models of the String group have appeared, each having different desirable features (see e.g. the appendix of [7] for seven such models). For instance, in one model [20], $\text{String}(n)$ is constructed as an extension as in eq. (1.2) of Lie groups so that $\widehat{G}$ has a Fréchet-Lie group structure uniquely determined up to isomorphism. The presence of such structures allows us to talk about $\text{String}(n)$ -principal bundles and String manifolds, regarding $\text{String}(n)$ as a structure group.

Studying the Spin group and its classifying space from the cohomology point of view requires understanding of the first generator, i.e. the generator in degree three the cohomology of the Spin group or the generator in degree four of the corresponding classifying space (see [36] [7] for interesting relations between the two). It is known from [32] that the cohomology ring of $B\text{Spin}$ in the stable case is generated by the Spin characteristic classes, the degree four generator of which is $\frac{1}{2}p_1$. However, we are interested in the unstable case. From the calculation presented by Kono [15] and Benson and Wood [3], we will deduce that $\frac{1}{2}p_1$ is also the generator of the cohomology $H^4(B\text{Spin}(n); \mathbb{Z})$. This is certainly known to experts working in this area, but we have not found it proved explicitly in the literature, and hence we include it for completeness and to ensure that we are on solid ground when discussing the unstable indefinite case. The lifting of the structure group from $\text{Spin}(n)$ to $\text{String}(n)$ of a bundle over a manifold X is then possible when the obstruction $\frac{1}{2}p_1(X)$ vanishes. We will be interested in generalizing this result to the case of $B\text{Spin}(p, q)$.

String structures are interesting from the geometric point of view due to the relation between the Riemannian geometry of a manifold and characteristic classes associated with String structures on that manifold. The Stolz-Höhn conjecture says the following: Let X be a smooth closed string manifold of dimension $4k$. If X admits a Riemannian metric with positive Ricci curvature, then the Witten genus $\phi_W(X)$ vanishes.

Homotopy-theoretically, String structures amount to having a String orientation, which in relation to modular forms gives a corresponding orientation to the spectrum of topological modular form (tmf), $M\text{String} \rightarrow \text{tmf}$ [1]. Conjecturally, the Witten genus is constructed as an index of a Dirac operator on loop space [40]. This is the String analog to the theorem on vanishing $\hat{A}$ -genus by Lichnerowicz [17] for Spin manifolds: Let X be a smooth closed spin manifold of dimension $2k$. If X admits a Riemannian affine connection with nonnegative and not identically zero Riemannian curvature, then the $\hat{A}$ -genus vanishes.

The Atiyah-Bott-Shapiro map $M\text{Spin} \rightarrow KO$ is constructed using the representations of the Spin groups, and relies on knowing that for a space X , elements of $KO^0(X)$ are represented by vector bundles over X . It gives a KO -theory Thom isomorphism for Spin-vector bundles, and is topological expression for the index of the Dirac operator. We hope that similar questions, also as in the seminal paper [40] and the considerable subsequent research that build on it, can be explored in the semi-Riemannian setting.

This paper is organized as follows. We start with a more general setting of the problem that we hope explain some of the homotopy theoretical constructions encountered here as well as in previous literature. In Sec. 2.1 and 2.2, we describe the Postnikov tower and the Whitehead tower of a space in a manner that is appropriate for applications. Then in Sec. 2.3 we provide a variant point of view on the Whitehead tower in way of clarification. Since lifts of indefinite Lie groups will be determined by their maximal compact subgroups, which are products, we discuss in Sec. 2.4 useful conditions on behavior of cohomology of products. The discussion is needed as our coefficients are the integers, which are not a field. This is applied in Sec. 2.5 to classifying spaces, where we identify the obstructions and where the fibrations become fiber bundles. We start with the applications in Sec. 3. First, in order to make sure we are on firm ground, we discuss the indefinite orthogonal groups in Sec. 3.1, highlighting their unstable homotopy groups. A subtle issue in the non-vanishing of the fundamental group of the corresponding Spin groups is addressed in Sec. 3.2. This is used in Sec. 3.3 to define the indefinite String groups, where we identify the obstructions explicitly by studying the generators of the classifying space of $B\text{Spin}(n)$ in the unstable case. We then show in Sec. 3.4 how the definitions and constructions extend to the case of $U(p, q)$ and $\text{Sp}(p, q)$, where subtle issues with stability are absent. Finally, in Sec. 3.5 we describe relations to twisted structures, and end with highlighting future work that we hope to do.

2 The Postnikov tower, Whitehead tower, and variants

In this section we will provide a careful treatment of the towers arising in co-killing homotopy groups of Lie groups. The idea is that Postnikov towers arise when killing homotopy groups above a certain degree, while Whitehead towers arise when killing homotopy groups below a certain degree—hence the term co-killing. We will also provide a variant tower of higher connected covers, which we demonstrate is equivalent to the latter. We believe that such a treatment, while certainly known to experts, seems to be missing from existing literature to the best of our knowledge. This technical treatment, we hope, will be for the benefit of the reader and will make the paper self-contained. Readers not interested in these details might wish to skip this section.

2.1 The Postnikov tower

We start by recalling the Postnikov tower (see [12] [11]). We will take this as our starting point to connect to the Whitehead tower.

Theorem 2.1 (see [12] [11]) *Let X be a connected space with a map $\alpha : X \rightarrow X\langle 1 \rangle$, where $X\langle 1 \rangle$ is the Eilenberg-Mac Lane space $K(\pi_1(X), 1)$, that induces an isomorphism on fundamental groups $\pi_1(X) \rightarrow \pi_1(X\langle 1 \rangle)$. Then there are spaces $X\langle n \rangle$ with maps $\alpha_n : X \rightarrow X\langle n \rangle$ which induce isomorphisms on homotopy groups $\pi_k(X) \rightarrow \pi_k(X\langle n \rangle)$ for $k \leq n$ and $\pi_k(X\langle n \rangle) = 0$ for $k > n$ with fibrations $p_{n+1} : X\langle n+1 \rangle \rightarrow X_n$ such that $\alpha_n = p_{n+1} \circ \alpha_{n+1}$.*

Assuming that such $\alpha_n : X \rightarrow X\langle n \rangle$ is given, the space $X\langle n+1 \rangle$ will be defined as the homotopy fiber of a certain map $k^{n+2} : X\langle n \rangle \rightarrow K(\pi_{n+1}(X), n+2)$ that is equivalent to the corresponding cohomology class in $H^{n+2}(X\langle n \rangle, \pi_{n+1}(X))$ such that the induced class $\alpha_n^* k^{n+2} \in H^{n+2}(X, \pi_{n+1}(X))$ is trivial. By the universality of the homotopy fiber $X\langle n+1 \rangle$, one then obtain the desired map $\alpha_{n+1} : X \rightarrow X\langle n+1 \rangle$. These maps are assembled in the following diagram called the *Postnikov*

tower of X :

$$\begin{array}{ccc}
 & \vdots & \\
 & \downarrow & \\
 & X\langle n \rangle \xrightarrow{k^{n+2}} K(\pi_{n+1}(X), n+2) & \\
 & \downarrow p_n & \\
 & \vdots & \\
 & \downarrow p_3 & \\
 & X\langle 2 \rangle \xrightarrow{k^4} K(\pi_3(X), 4) & \\
 & \downarrow p_2 & \\
 X & \xrightarrow{\alpha_1} X\langle 1 \rangle \xrightarrow{k^3} K(\pi_2(X), 3) & \\
 & \nearrow \alpha_2 & \\
 & \nearrow \alpha_n &
 \end{array}
 \tag{2.1}$$

One could see how to choose such a map $k^{n+2} : X\langle n \rangle \rightarrow K(\pi_{n+1}(X), n+2)$ and how to obtain $X\langle n+1 \rangle$ as the homotopy fiber explicitly as follows. Let $j : C(\alpha_n) \rightarrow K(\pi_{n+1}(X), n+2)$ be the inductive attachment of cells to the cofiber $C(\alpha_n)$ to match the homotopy group of the space $K(\pi_{n+1}(X), n+2)$ from the identification $\pi_{n+2}(C(\alpha_n)) \cong \pi_{n+1}(X)$ and the fact that $C(\alpha_n)$ is $(n+1)$ -connected. On the other hand, we have the inclusion $X\langle n \rangle \hookrightarrow C(\alpha_n)$ and we take the composite $k^{n+2} : X\langle n \rangle \rightarrow K(\pi_{n+1}(X), n+2)$. From the inclusion of the cone $C(X)$ in the cofiber $C(\alpha_n)$, $\chi_x(t) := j(x, 1-t)$ for $(x, 1-t) \in C(X)$ is in the path space $PK(\pi_{n+1}(X), n+2)$. This then makes the outer rectangle part of the following pullback diagram

$$\begin{array}{ccc}
 X & \xrightarrow{x \mapsto \chi_x} & PK(\pi_{n+1}(X), n+2) \\
 \searrow \alpha_{n+1} & & \downarrow \text{ev}_1 \\
 X\langle n+1 \rangle & \longrightarrow & PK(\pi_{n+1}(X), n+2) \\
 \downarrow p_{n+1} & & \downarrow \text{ev}_1 \\
 X\langle n \rangle & \xrightarrow{k^{n+2}} & K(\pi_{n+1}(X), n+2)
 \end{array}
 \tag{2.2}$$

commutative, so that we have the homotopy fiber $X\langle n+1 \rangle$. Here ev_1 denotes evaluation at the point 1 of the interval in the path space. Note that $PK(\pi_{n+1}(X), n+2)$ is a homotopy equivalent to a point space so the commutativity of the diagram implies that the cohomology class $\alpha_n^* k^{n+2} \in H^{n+2}(X; \pi_{n+1}(X))$ is trivial. Therefore, we obtain the map α_{n+1} which satisfies $\alpha_n = p_{n+1} \circ \alpha_{n+1}$ and induces an isomorphism $\pi_k(X) \rightarrow \pi_k(X\langle n+1 \rangle)$ for $k \leq n+1$ and makes $X\langle n+1 \rangle$ have trivial k^{th} homotopy group for $k > n+2$. By inductive application of this process, one obtains the *Postnikov tower* of X .

2.2 The Whitehead tower

Next we consider the Whitehead tower, which is in a sense a dual to the Postnikov tower [38] [39]. Note that the Postnikov tower approximates the homotopy groups of X “from the bottom” in the sense that it admits an isomorphism of lower homotopy groups with higher homotopy groups being killed. One then may seek the dual process of estimating the homotopy groups of X “from the top”,

in the sense that lower homotopy groups are killed while higher homotopy groups are isomorphic to those of X .

Theorem 2.2 (see [39]) *Let X be a connected space with a Postnikov tower. Then there are spaces $X\langle n \rangle$ such that $\pi_k(X\langle n \rangle) = 0$ for $k \leq n$ and maps $\hat{\alpha}_n : X\langle n \rangle \rightarrow X$ that induce isomorphisms $\pi_k(X\langle n \rangle) \rightarrow \pi_k(X)$ for $k > n$. Moreover, there is a fibration $\hat{p}_{n+1} : X\langle n+1 \rangle \rightarrow X\langle n \rangle$ for each n such that $\hat{\alpha}_{n+1} = \hat{\alpha}_n \circ \hat{p}_{n+1}$ with a fiber the based loop space $\Omega X\langle n \rangle$.*

From the Postnikov tower of X , we have spaces $X\langle n \rangle$ and maps $\alpha_n : X \rightarrow X\langle n \rangle$. By taking the homotopy fiber $X\langle n \rangle$ of $\alpha_n : X \rightarrow X\langle n \rangle$, we obtain a fibration $\hat{\alpha}_n : X\langle n \rangle \rightarrow X$ that induces isomorphisms $\pi_k(X\langle n \rangle) \rightarrow \pi_k(X)$ for $k > n$ and makes $\pi_k(X\langle n \rangle) = 0$ for $k \leq n$. Since $X\langle n+1 \rangle \rightarrow X$ factors through $X\langle n \rangle$ naturally as in the diagram

$$\begin{array}{ccccc}
 \hat{X}\langle n+1 \rangle & \xrightarrow{\quad} & PX\langle n+1 \rangle & & \\
 \searrow \hat{p}_{n+1} & & \searrow P(p_{n+1}) & & \\
 & & X\langle n \rangle & \xrightarrow{\quad} & PX\langle n \rangle \\
 \searrow \hat{\alpha}_{n+1} & & \downarrow \hat{\alpha}_n & & \downarrow \\
 & & X & \xrightarrow{\alpha_n} & X\langle n \rangle
 \end{array} \tag{2.3}$$

we obtain a map $\hat{p}_{n+1} : \hat{X}\langle n+1 \rangle \rightarrow X\langle n \rangle$, which can be made into a fibration, up to homotopy equivalence, such that $\hat{\alpha}_{n+1} = \hat{\alpha}_n \circ \hat{p}_{n+1}$. The induced long exact sequence of the fibration $\hat{p}_{n+1}$ shows that the fiber is homotopy equivalent to the Eilenberg-MacLane space $K(\pi_{n+1}(X), n)$. The tower of fibrations thus obtained is a dual version of the Postnikov tower called the *Whitehead tower* [38] [39]:

$$\begin{array}{ccc}
 \vdots & & \\
 \downarrow & & \\
 K(\pi_n(X), n-1) \longrightarrow & X\langle n \rangle & \\
 & \downarrow \hat{p}_n & \\
 \vdots & & \\
 K(\pi_3(X), 2) \longrightarrow & X\langle 3 \rangle & \\
 & \downarrow \hat{p}_3 & \\
 K(\pi_2(X), 1) \longrightarrow & X\langle 2 \rangle & \\
 & \downarrow \hat{p}_2 & \\
 & X\langle 1 \rangle & \\
 & \xrightarrow{\hat{\alpha}_1} & X.
 \end{array}
 \tag{2.4}$$

$\hat{\alpha}_n$
 $\hat{\alpha}_3$
 $\hat{\alpha}_2$

2.3 A variant view

Each fibration $\widehat{p}_{n+1} : X\langle n+1 \rangle \rightarrow X\langle n \rangle$ in the Whitehead tower can be regarded as the $(n+1)$ -connected covering of $X\langle n \rangle$ and we may ask the “lifting” question, i.e. under what circumstance can a map $M \rightarrow X\langle n \rangle$ be lifted to $M \rightarrow X\langle n+1 \rangle$ as in the commutative diagram

$$\begin{array}{ccc} & & X\langle n+1 \rangle \\ & \nearrow \text{dashed} & \downarrow \\ M & \longrightarrow & X\langle n \rangle . \end{array} \quad (2.5)$$

To answer this question we need the following important property of the sequence of spaces appearing in the Whitehead tower.

Proposition 2.3 *Let X be a connected space. Then there are spaces $X\langle n \rangle$, where $X\langle 1 \rangle := X$, with fibrations $q_{n+1} : X\langle n+1 \rangle \rightarrow X\langle n \rangle$ with fiber $K(\pi_n(X), n-1)$ such that*

$$\pi_k(X\langle n \rangle) \cong \begin{cases} 0 & \text{for } k < n \\ \pi_k(X) & \text{for } k \geq n. \end{cases}$$

Remark 1 *These spaces $X\langle n+1 \rangle$ are constructed as the homotopy fibers of a cohomology class $\lambda_n \in H^n(X\langle n \rangle; \pi_n(X))$ that induces an isomorphism $\pi_n(X\langle n \rangle) \xrightarrow{\cong} \pi_n(X)$, and a map $f : M \rightarrow X\langle n \rangle$ is lifted to $\widetilde{f} : M \rightarrow X\langle n+1 \rangle$ if the induced cohomology class $f^*\lambda_n \in H^n(M; \pi_n(X))$ is trivial.*

Thus we have the following diagram that one might call the *tower of higher connected covers*

$$\begin{array}{ccccc} & & \vdots & & \\ & & \downarrow & & \\ & & X\langle n \rangle & \xrightarrow{\lambda_n} & K(\pi_n(X), n) \\ & \nearrow \text{dashed} & \downarrow q_n & & \\ & & \vdots & & \\ & & \downarrow q_3 & & \\ & & X\langle 2 \rangle & \xrightarrow{\lambda_2} & K(\pi_2(X), 2) \\ & \nearrow & \downarrow \lambda_2 & & \\ M & \longrightarrow & X\langle 1 \rangle & \xrightarrow{\lambda_1} & K(\pi_1(X), 1) . \end{array} \quad (2.6)$$

The construction is an inductive application of the process in the following lemma.

Lemma 2.4 (Killing n^{th} homotopy group) *Let X be a simply connected space and Y be a space such that $\pi_{n-1}(Y) = 0$ and the cohomology class $\lambda_n \in H^n(Y; \pi_n(X))$ induces an isomorphism $\pi_n(Y) \rightarrow \pi_n(X)$. Then the homotopy fiber $\widehat{Y}$ of λ_n satisfies*

$$\pi_k(\widehat{Y}) \cong \begin{cases} 0 & \text{for } k = n \\ \pi_k(Y) & \text{for } k \neq n, \end{cases}$$

with a fibration $q : \widehat{Y} \rightarrow Y$. A map $f : M \rightarrow Y$ is lifted to $\widehat{f} : M \rightarrow \widehat{Y}$ with respect to the fibration q when the induced cohomology class $f^*\lambda_n \in H^n(M; \pi_n(X))$ is trivial.

Proof. We would like to consider the lifting

$$\begin{array}{ccc} & \widehat{Y} & \\ \widehat{f} \nearrow & \downarrow q & \\ M & \xrightarrow{f} Y & \xrightarrow{\lambda_n} K(\pi_n(X), n) \end{array} \quad (2.7)$$

The result then follows from considering the following pullback diagram

$$\begin{array}{ccccc} M & & & & \\ & \searrow \widehat{f} & & \searrow & \\ & \widehat{Y} & \xrightarrow{\quad} & PK(\pi_n(X), n) & \\ & \downarrow q \lrcorner & & \downarrow \text{ev}_1 & \\ & Y & \xrightarrow{\lambda_n} & K(\pi_n(X), n) & \end{array} \quad (2.8)$$

and its long exact sequence of homotopy groups and the fact that the path space $PK(\pi_n(X), n)$ is contractible. Here, the map ev_1 is the evaluation map at the point 1 in the path. $\square$

Remark 2 (i) Although the fibrations $\widehat{p}_n : \widehat{X}_n \rightarrow \widehat{X}_{n-1}$ and $q_{n+1} : X\langle n+1 \rangle \rightarrow X\langle n \rangle$ may look similar by having the same fiber $K(\pi_n(X), n-1)$, a comparison is needed. One way of figuring out that a given fibration $K(\pi_n(X), n-1) \rightarrow X\langle n+1 \rangle \rightarrow X\langle n \rangle$ fits in the tower of higher connected covers is to check that a map $f : M \rightarrow X\langle n \rangle$ can be lifted up to a map $\widehat{f} : M \rightarrow X\langle n+1 \rangle$ if and only if cohomology class $f^*\lambda_n \in H^n(M; \pi_n(X))$ induced by λ_n which induced the isomorphism of n^{th} -homotopy group vanishes.

(ii) It is convenient to rework the definition in a way making explicit the obstruction problem giving the lift from $X\langle n-1 \rangle$ to $X\langle n \rangle$. Indeed, the two constructions, namely the Whitehead tower and the tower of higher connected covers, are equivalent by the homotopy commutative diagram:

$$\begin{array}{ccccc} X\langle n \rangle & \xrightarrow{\quad} & * & & \\ \downarrow & & \downarrow & & \\ X\langle n-1 \rangle & \xrightarrow{\quad} & K(\pi_n(X), n) & \xrightarrow{\quad} & * \\ \downarrow & & \downarrow & & \downarrow \\ X & \xrightarrow{\quad} & X\langle n \rangle & \xrightarrow{\quad} & X\langle n-1 \rangle \end{array} \quad (2.9)$$

2.4 Computational aspects

We now turn to useful computational aspects of the above constructions. This will also be useful when considering indefinite Lie groups in Sec. 3. In the notation of the previous section, let us assume that $H^n(Y; \pi_n(X))$ is computable and let us see which of its classes induce isomorphisms

$\pi_n(Y) \cong \pi_n(X)$. If $H^n(Y; \pi_n(X))$ is generated by a single class α , then the map $\alpha : Y \rightarrow K(\pi_n(X), n)$ induces an isomorphism $\pi_n(Y) \xrightarrow{\cong} \pi_n(X)$. Indeed, there is an isomorphism in the bracket $[\pi_n(Y), \pi_n(X)]$ that corresponds to $n\alpha$ for some integer n , so the generator α has to induce an isomorphism.

Suppose that n th homotopy group of Y is of the form $A \times B$ for some abelian groups A and B such that each of $H^n(Y; A)$ and $H^n(Y; B)$ has a single generator α and β , respectively. Then α and β induce group homomorphism $\pi_n(Y) \rightarrow A$ and $\pi_n(Y) \rightarrow B$ so that neither $(\alpha, 0)$ nor $(0, \beta)$ in $H^n(Y; A \times B)$ can induce an isomorphism $\pi_n(Y) \xrightarrow{\cong} A \times B$. However, the class in $H^n(Y; A) \times H^n(Y; B)$ inducing the isomorphism does exist and is of the form $(\tilde{\alpha}, \tilde{\beta})$ where $\tilde{\alpha} = a\alpha \in H^n(Y; A)$ and $\tilde{\beta} = b\beta \in H^n(Y; B)$ for some nonzero integers a and b .

If Y is homotopy equivalent to the product space $Y' \times Y''$ and

$$H^n(Y; \pi_n(X)) \cong H^n(Y'; \pi_n(X)) \times H^n(Y''; \pi_n(X)) \quad (2.10)$$

then we have the decomposition

$$H^n(Y; \pi_n(X)) \cong H^n(Y'; \pi_n(Y')) \times H^n(Y'; \pi_n(Y'')) \times H^n(Y''; \pi_n(Y')) \times H^n(Y''; \pi_n(Y'')) . \quad (2.11)$$

We would like to know when this occurs. A direct check leads to the following

Lemma 2.5 *A product space $Y' \times Y''$ satisfies the condition of eq. (2.10) if all of the following are satisfied:*

1. *The chain complex for Y' is flat;*
2. $\bigoplus_{p+q=n} H_p(Y'; \mathbb{Z}) \times H_q(Y''; \mathbb{Z}) \cong H_n(Y'; \mathbb{Z}) \oplus H_n(Y''; \mathbb{Z});$
3. $\bigoplus_{p+q=n-1} \text{Tor}_1^{\mathbb{Z}}(H_p(Y'; \mathbb{Z}), H_q(Y''; \mathbb{Z})) = 0;$
4. $\text{Ext}_{\mathbb{Z}}^1(H_{n-1}(W; \mathbb{Z}), \pi_n(X)) = 0$ for $W = Y', Y''$ and $Y' \times Y''$.

If conditions 1, 2 and 3 hold, then the Künneth theorem gives an isomorphism

$$H_n(Y' \times Y''; \mathbb{Z}) \cong H_n(Y'; \mathbb{Z}) \oplus H_n(Y''; \mathbb{Z}) . \quad (2.12)$$

If, in addition, condition 4 also holds, then the universal coefficient theorem implies, for $W = Y', Y''$ and $Y' \times Y''$, that

$$\text{hom}(H_n(W; \mathbb{Z}), \pi_n(X)) \cong H^n(W; \pi_n(X)) , \quad (2.13)$$

which gives the isomorphism of eq. (2.10).

Assuming further that both $H^n(Y'; \pi_n(Y'))$ and $H^n(Y''; \pi_n(Y''))$ are generated by single classes α' and α'' respectively, the cohomology class (α', α'') induces the desired isomorphism $\pi_n(Y' \times Y'') \xrightarrow{\cong} \pi_n(Y') \times \pi_n(Y'')$. Obviously, the cohomology class (α', α'') is the same as $\alpha' \times \alpha'' : Y' \times Y'' \rightarrow K(\pi_n(Y'), n) \times K(\pi_n(Y''), n)$. Thus, we have obtained:

Proposition 2.6 *For a space Y such that $\pi_{n-1}(Y) = 0$ and $\pi_n(Y) \cong \pi_n(X)$, the homotopy fiber $\hat{Y}$ of $\lambda_n : Y \rightarrow K(\pi_n(X), n)$ has homotopy groups isomorphic to Y except $\pi_n(\hat{Y}) = 0$, according to the following cases:*

(i) $\lambda_n = \alpha$ if α is the only generator of $H^n(Y, \pi_n(X))$, i.e.

$$\begin{array}{ccc} \widehat{Y} & \xrightarrow{\quad} & \text{pt} \\ \downarrow \lrcorner & & \downarrow \\ Y & \xrightarrow{\alpha} & K(\pi_n(X), n) . \end{array}$$

(ii) $\lambda_n = (\widehat{\alpha}, \widehat{\beta})$ if $\pi_n(Y) \cong A \times B$ and α and β are the only generators of $H^n(Y; A)$ and $H^n(Y; B)$, respectively, where $\widehat{\alpha} = a\alpha$ and $\widehat{\beta} = b\beta$ for some nonzero integers a and b , respectively, i.e.

$$\begin{array}{ccc} \widehat{Y} & \xrightarrow{\quad} & \text{pt} \\ \downarrow \lrcorner & & \downarrow \\ Y & \xrightarrow{(\widehat{\alpha}, \widehat{\beta})} & K(A \times B, n) . \end{array}$$

(iii) $\lambda_n = \alpha' \times \alpha''$ which is equivalent to the map $Y' \times Y'' \rightarrow K(\pi_n(Y'), n) \times K(\pi_n(Y''), n)$ if $Y \simeq Y' \times Y''$ where α', α'' are the only generators of $H^n(Y', \pi_n(Y'))$ and $H^n(Y'', \pi_n(Y''))$ respectively, and $H^n(Y; \pi_n(X))$ is decomposed as in eq. (2.10), i.e.

$$\begin{array}{ccc} \widehat{Y} & \xrightarrow{\quad} & \text{pt} \\ \downarrow \lrcorner & & \downarrow \\ Y' \times Y'' & \xrightarrow{\alpha' \times \alpha''} & K(\pi_n(Y') \times \pi_n(Y''), n) . \end{array}$$

2.5 Special case: higher connected covers of classifying spaces and obstructions

We will mainly be interested in our spaces being topological groups. Let G be a topological group with a trivial particular homotopy group $\pi_{n-1}(G) = 0$. Then BG is a classifying space of G such that $\pi_n(BG) = 0$. By taking the homotopy fiber of some map $BG \rightarrow K(\pi_{n-1}(BG), n-1)$, we obtain a space $\widehat{BG}$ for which $\pi_{n+1}(\widehat{BG})$ is trivial. Then its loop space $\Omega\widehat{BG}$ satisfies $\pi_n(\Omega\widehat{BG}) = 0$. Setting $\widehat{G} := \Omega\widehat{BG}$, we get a topological space which is G with π_n killed. The immediate consequence of this observation is the following:

Lemma 2.7 $\widehat{BG}$ is homotopy equivalent to $B(\widehat{G})$.

We now consider classification of the corresponding bundles. From the general discussion in previous sections we have the following results.

Proposition 2.8 Suppose that there is a G -principal bundle over M determined by a classifying map $f : M \rightarrow BG$. Then there exists a map $\widehat{f} : M \rightarrow B\widehat{G}$ providing a $\widehat{G}$ -principal bundle over the same base M compatible with the bundle f with respect to the map $B\widehat{G} \rightarrow BG$ if $f^*\lambda_n \in H^{n+1}(M, \pi_n(G_{n-1}))$ is trivial for some $\lambda_n \in H^n(BG; \pi_{n-1}(G))$ that induces an isomorphism $\pi_n(BG) \rightarrow \pi_{n-1}(G)$.

Remark 3 (i) *The condition in Proposition (2.8) is equivalent to saying that $f^*\lambda_n$ factors through a point space up to homotopy:*

$$\begin{array}{ccc}
 M & \xrightarrow{\quad \quad} & \text{pt} \\
 \downarrow f & \nearrow \hat{f} & \downarrow \\
 B\hat{G} & \xrightarrow{\quad \quad} & \text{pt} \\
 \downarrow & & \downarrow \\
 BG & \xrightarrow{\lambda_n} & K(\pi_{n-1}(G), n) .
 \end{array} \tag{2.14}$$

(ii) *We say that the obstruction $f^*\lambda_n$ vanishes or trivialized so that the bundle f is lifted to $\hat{f}$. This lifting of f to $\hat{f}$ is also called the trivialization of the class λ_n .*

Indeed, the pullback diagram eq. (2.8) induces another pullback diagram classifying $\hat{G}$ -principal bundles over X . Denoting the set of G -principal bundles over X by $\text{Bun}_G(X)$, we have a pullback diagram

$$\begin{array}{ccc}
 \text{Bun}_{\hat{G}}(X) & \xrightarrow{\quad \quad} & \text{pt} \\
 \downarrow \lrcorner & & \downarrow \\
 \text{Bun}_G(X) & \longrightarrow & H^n(X; \pi_{n-1}(G)) .
 \end{array} \tag{2.15}$$

Inductively, we get a tower of connected covers which is the right side of the diagram:

$$\begin{array}{ccc}
 \vdots & & \vdots \\
 \downarrow & & \downarrow \\
 BG\langle n \rangle & \xrightarrow{\lambda_n} & K(\pi_{n-1}(G), n) \\
 \downarrow & & \downarrow \\
 \vdots & & \vdots \\
 \downarrow q_2 & & \downarrow \\
 BG\langle 2 \rangle & \xrightarrow{\lambda_2} & K(\pi_1(G), 2) \\
 \downarrow q_1 & & \downarrow \\
 BG\langle 1 \rangle & \xrightarrow{\lambda_1} & K(\pi_0(G), 1) \\
 \downarrow q_0 & & \downarrow \\
 X & \xrightarrow{\quad \quad} & BG \xrightarrow{\lambda_0} K(\pi_0(BG), 0) .
 \end{array} \tag{2.16}$$

More particularly, we will be mainly interested in the case when $G\langle n \rangle$ is an $(n-1)$ -connected topological group, and also in the classifying space BG for some topological group G , i.e., $X = BG$, and its $(n-1)$ -connected cover $BG\langle n \rangle$. We also assume, without loss of generality, that $\pi_{n+1}(BG\langle n \rangle) \neq 0$ and our goal would be to obtain a topological group $\hat{G}$ such that $\widehat{BG} \simeq B(\hat{G})$.

Starting with the latter, we have the homotopy pullback diagram

$$\begin{array}{ccc} \widehat{BG} & \longrightarrow & \text{pt} \\ \downarrow & & \downarrow \\ BG & \longrightarrow & K(\pi_n(G), n+1) \end{array} \quad (2.17)$$

and the fibration

$$K(\pi_n(G), n) \rightarrow \widehat{BG} \rightarrow BG . \quad (2.18)$$

Having a homotopy pullback diagram such as (2.17) is equivalent to saying that $\widehat{BG}$ is the homotopy fiber of $BG \rightarrow K(\pi_n(G), n+1)$, hence equivalent to saying that $\widehat{BG} \rightarrow BG \rightarrow K(\pi_n(G), n+1)$ is a homotopy fiber sequence.

Next, setting $\widehat{G} = \Omega \widehat{BG}$ we have $B\widehat{G} \simeq \widehat{BG}$, and this $\widehat{G}$ fits into the diagram

$$\begin{array}{ccc} \widehat{G} & \longrightarrow & \text{pt} \\ \downarrow & & \downarrow \\ G & \longrightarrow & K(\pi_n(G), n) , \end{array} \quad (2.19)$$

which induces a fibration with fiber $K(\pi_n(G), n-1)$

$$K(\pi_n(G), n-1) \rightarrow \widehat{G} \rightarrow G . \quad (2.20)$$

Since $K(\pi_n(G), n-1)$ has a group structure, then this induces (2.18) as a fiber bundle. In fact, this will be a *principal fiber bundle*, the point being that the homotopy fiber of a map to a connected space X is actually an ΩX -principal bundle with the structure group an actual topological group representing the homotopy type of the loop space ΩX . Furthermore, in that case, $\widehat{G}$ itself is directly defined to be the sequence in eq. (2.20) as a principal fiber bundle.

For example, for the case $n = 1$, we have $\widehat{G} = \text{Spin}(m)$ defined in terms of Clifford algebra to be the double cover of $G = \text{SO}(m)$, $m \geq 3$.

Let G be a topological group with $\pi_{n+1}(BG) \cong \pi_n(G) \neq 0$. Also, let $\widehat{G}$ be a topological group homotopy equivalent to the loop space $\Omega \widehat{BG}$ of $\widehat{BG}$. This can be viewed as the homotopy fiber of $\lambda_{n+1} : BG \rightarrow K(\pi_n(G), n+1)$ determined according to Proposition 2.6 so that $\pi_n(G)$ is killed, i.e., $\pi_i(\widehat{G}) \cong \pi_i(G)$ for $i \geq 0$ and $i \neq n$, and $\pi_n(\widehat{G}) = 0$. Then we obtain a fibration

$$K(\pi_n(G), n+1) \rightarrow B\widehat{G} \rightarrow BG . \quad (2.21)$$

On the other hand, suppose that we have a fibration $\widehat{G} \rightarrow G$. Then the homotopy groups of $\widehat{G}$ and G forces the kernel to be $K(\pi_n(G), n)$ so that we have a fibration (2.20), which induces the fibration (2.21) as $K(\pi_n(G), n)$ has the group structure.

Proposition 2.9 *The fibrations in the Whitehead tower (hence also the tower of connected covers) of a Lie group are principal fiber bundles.*

3 Applications

3.1 Special orthogonal groups $\mathrm{SO}(n)$ and $\mathrm{SO}(p, q)$

We now concentrate on the orthogonal group and its connected covers. We start with the definite signature and then work our way to indefinite signatures. Starting with the former, we have the statement that identifies when the shift in rank is inconsequential

$$\pi_i(\mathrm{O}(n)) \cong \pi_i(\mathrm{O}(n+1)) \quad \text{for } 0 < i \leq n-2. \quad (3.1)$$

Note that according to Kervaire [13], this cannot be improved in general as $\pi_6(\mathrm{SO}(7)) \cong 0$, while $\pi_7(\mathrm{SO}(7)) \cong \mathbb{Z}$. The following isomorphism shows that the special orthogonal group $\mathrm{SO}(n)$ has the same homotopy groups as the orthogonal group $\mathrm{O}(n)$ in positive degrees

$$\pi_i(\mathrm{SO}(n)) = \begin{cases} 0 & \text{for } i = 0 \\ \pi_i(\mathrm{O}(n)) & \text{for } i \geq 1. \end{cases} \quad (3.2)$$

We also note the following low degree identifications, which are often useful in calculations and applications: $\mathrm{O}(1) \cong S^0$, $\mathrm{SO}(2) \cong S^1$, $\mathrm{SO}(3) \cong \mathbb{R}P^3$, $\mathrm{SO}(4) \cong S^3 \times S^3$.

Collecting all the above observations and results gives the following table of *unstable* homotopy groups of lower dimensional orthogonal groups (see e.g. [13]):

	O(1)	O(2)	O(3)	O(4)	O(5)	O(6)	O(7)	O(8)	O(9)
π_0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$
π_1	0	$\mathbb{Z}$	$\mathbb{Z}/2$						
π_2	0	0	0	0					
π_3	0	0	$\mathbb{Z}$	$\mathbb{Z} \times \mathbb{Z}$	$\mathbb{Z}$				
π_4	0	0	$\mathbb{Z}/2$	$\mathbb{Z}/2 \times \mathbb{Z}/2$	$\mathbb{Z}/2$	0			
π_5	0	0	$\mathbb{Z}/2$	$\mathbb{Z}/2 \times \mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}$	0		
π_6	0	0	$\mathbb{Z}/12$	$\mathbb{Z}/12 \times \mathbb{Z}/12$	0	0	0	0	
π_7	0	0	$\mathbb{Z}/2$	$\mathbb{Z}/2 \times \mathbb{Z}/2$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z} \times \mathbb{Z}$	$\mathbb{Z}$

where the boxed entries indicate that the corresponding homotopy groups are being stabilized.

We now consider the corresponding classifying spaces. We know that the first cohomology $H^1(\mathrm{BO}(n), \mathbb{Z}/2) \cong \mathbb{Z}/2$ is generated by the first Stiefel-Whitney class w_1 , so that we may pullback $w_1 : \mathrm{BO}(n) \rightarrow K(\mathbb{Z}/2, 1)$ to obtain a 0-connected cover, say G , of $\mathrm{O}(n)$:

$$\begin{array}{ccccc} & & BG & \longrightarrow & \mathrm{pt} \\ & \nearrow \hat{f} & \downarrow & & \downarrow \\ X & \xrightarrow{f} & \mathrm{BO}(n) & \xrightarrow{w_1} & K(\mathbb{Z}/2, 1). \end{array} \quad (3.3)$$

We also know that the special orthogonal group $\mathrm{SO}(n)$ has the same homotopy groups as G so we may regard the connected identity component $\mathrm{SO}(n)$ of $\mathrm{O}(n)$ as the 0-connected cover G . The pattern continues to the Spin group and beyond, as in [27].

Next, consider our main object which is the indefinite orthogonal group $O(p, q)$. This has a maximal compact subgroup $O(p) \times O(q)$. The inclusion $SO(p) \times SO(q) \hookrightarrow SO(p, q)$ is a homotopy equivalence by the Cartan decomposition of noncompact Lie groups, i.e. the homeomorphism from $SO(p, q) \rightarrow SO(p) \times SO(q) \times \mathbb{R}^{pq}$.

With this observation, we can reduce the problem of connected covers in this indefinite signature setting to essentially two copies of the problem in the definite signature case. From the results in the previous section, we may pullback $w_1 \times w_1 : BO(p) \times BO(q) \rightarrow K(\mathbb{Z}/2, 1) \times K(\mathbb{Z}/2, 1)$ to obtain the 0-connected cover G :

$$\begin{array}{ccccc} & BG(p, q) & \longrightarrow & \text{pt} & \\ & \downarrow & & \downarrow & \\ X & \xrightarrow{\hat{f}} & BO(p, q) & \xrightarrow{w_1 \times w_1} & K(\mathbb{Z}/2 \times \mathbb{Z}/2, 1) . \end{array} \quad (3.4)$$

We have $G(p, q) \simeq SO(p) \times SO(q)$ and this is homotopy equivalent to the identity component $SO(p, q)^0$ of $SO(p, q)$, so we may also take $G = SO(p, q)^0$ and denote this by $\widehat{SO}(p, q)$ for notational consistency. Note that $SO(1) \simeq \text{pt}$ and $\widehat{SO}(1, n) \simeq SO(n)$ so that we do not have to consider the case when $n = 1$ in killing its higher homotopy groups.

On the other hand, we have the following.

Definition 3.1 *The twisted covering, denoted by $\widetilde{SO}(p, q)$, is the pullback:*

$$\begin{array}{ccc} \widetilde{BSO}(p, q) & \longrightarrow & BO(q) \\ \downarrow \lrcorner & & \downarrow w_1 \\ BO(p) & \xrightarrow{w_1} & K(\mathbb{Z}/2, 1). \end{array} \quad (3.5)$$

3.2 Indefinite Spin groups

The next step in going from $SO(n)$ to $\text{Spin}(n)$ by taking a double cover, or killing the fundamental group, leads to the isomorphism

$$\pi_i(\text{Spin}(n)) = \begin{cases} 0 & \text{for } i \geq 2 \\ \pi_i(O(n)) & \text{for } i \geq 3. \end{cases} \quad (3.6)$$

Note the following useful isomorphisms in low degrees: $\text{Spin}(2) \cong U(1)$, $\text{Spin}(3) \cong \text{SU}(2)$, $\text{Spin}(5) \cong \text{Sp}(2)$, and $\text{Spin}(6) \cong \text{SU}(4)$.

We next consider the indefinite orthogonal group $SO(p, q)$ and kill the first homotopy group π_1 . Here there are two case, depending on whether one of the factors p or q is greater than 1. Thus we would like to kill the fundamental group of either $SO(1, n) \simeq O(n)\langle 1 \rangle \simeq SO(n)$ or $SO(p, q) \simeq O(p, q)\langle 1 \rangle$ for $p, q \geq 2$. An important distinction with the definite case is that the maximal compact subgroup of $SO(p, q)$ is $SO(p) \times SO(q)$, signalling that connectedness involves more than just the usual $\mathbb{Z}/2$.

Remark 4 (i). The group $\text{Spin}(p, q)$ is defined to be the double cover of $\text{SO}(p, q)$ in a slightly nontrivial way, namely the cover corresponding to the diagonal $\mathbb{Z}/2$ inside $\mathbb{Z}/2 \times \mathbb{Z}/2$. This diagonal $\mathbb{Z}/2$ is formed of a pair of elements, where the first correspond to the kernel of $\text{Spin}(p) \rightarrow \text{SO}(p)$ and the second to $\text{Spin}(q) \rightarrow \text{SO}(q)$. For example, $\text{Spin}(2, 2) \cong \text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R})$. See [41] [2] [16] [33] [34].

(ii). The condition for having a $\text{Spin}(p, q)$ structure from a $\text{SO}(p, q)$ structure is the separate vanishing of two second Stiefel-Whitney classes w_2^i , $i = 1, 2$ (see [29] for details).

Rationally, the cohomology ring of the special orthogonal group is given as

$$H^*(BSO(n); \mathbb{Q}) \cong \begin{cases} \mathbb{Q}[p_1, p_2, \dots, p_{\lfloor \frac{n}{2} \rfloor}], & n \text{ odd}, \\ \mathbb{Q}[p_1, p_2, \dots, p_{\frac{n}{2}}, e] / (p_{\frac{n}{2}} - e^2), & n \text{ even}, \end{cases} \quad (3.7)$$

where p_i are the Pontrjagin classes in degree $4i$ and e is the Euler class in degree n . The result on the odd case is what one expects, while the even case introduces a new generator. When considering the integral case, this generator persists and, in addition, we will have other generators arising from integral lifts of the Stiefel-Whitney classes, i.e. arising from applying the Bockstein on monomials in even Stiefel-Whitney classes. Since we are interested in degree four generators, the latter will not be of relevance to us.

We know that $H^2(BSO(n); \mathbb{Z}/2) \cong \mathbb{Z}/2$ with the second Stiefel-Whitney class w_2 as its generator. For $n = 2$, the integral cohomology of $BSO(2)$ is isomorphic to $\mathbb{Z}$ with a single generator $^1 \sqrt{p_1}$ such that $\sqrt{p_1} \sqrt{p_1} = p_1 \in H^4(BSO(2); \mathbb{Z})$ by the result of Brown on the integral cohomology ring of $BSO(n)$ [5]. In general, in the integral cohomology of $BSO(n)$ the square e^2 of the Euler class is the same as the Pontrjagin class in degree $4n$. So for $n = 1$, we have a generator of degree 2 given by $e = \sqrt{p_1}$. One can also view this as a first Chern class if one identifies $\text{SO}(2)$ with $\text{U}(1)$ and hence $BSO(n)$ with $\mathbb{C}P^\infty$, whose cohomology is given as $H^*(\mathbb{C}P^\infty; \mathbb{Z}) \cong \mathbb{Z}[x]$, with $|x| = 2$.

Therefore, we obtain a 1-connected cover $G(n)$ of $\text{SO}(n)$ for $n \geq 3$, and $G(2)$ of $\text{SO}(2)$, respectively, by taking pullbacks:

$$\begin{array}{ccc} BG(n) & \longrightarrow & \text{pt} \\ \nearrow & \downarrow & \downarrow \\ X & \longrightarrow BSO(n) \xrightarrow{w_2} & K(\mathbb{Z}/2, 2) \end{array}, \quad \begin{array}{ccc} BG(2) & \longrightarrow & \text{pt} \\ \nearrow & \downarrow & \downarrow \\ X & \longrightarrow BSO(2) \xrightarrow{\sqrt{p_1}} & K(\mathbb{Z}, 2) \end{array}. \quad (3.8)$$

We know that the spin group $\text{Spin}(n)$ is homotopy equivalent to $G(n)$ for $n \geq 3$. However, $\text{Spin}(2)$ and $G(2)$ do not agree since $\text{Spin}(2) \cong S^1$ is not simply connected.

Definition 3.2 The homotopic (definite) spin group, denoted by $\widehat{\text{Spin}}(n)$, is the 1-connected cover of $\text{SO}(n)$ for all $n \geq 2$ so that $\widehat{\text{Spin}}(n) \simeq \text{Spin}(n)$ and $\widehat{\text{Spin}}(2) \simeq G(2)$.

In fact, as far as killing higher homotopy groups is concerned, $G(2)$ is trivial and we do not have to consider the case with $n = 2$ in the process.

Next, in order to kill π_1 of $\text{SO}(p, q)$, we consider three cases: When $p = q = 2$, when either p or q is equal to 2, and when both p and q are greater than 2.

¹Here we are calling the generators $\sqrt{p_1}$ as these square to the first Pontrjagin class. At the level of $\mathbb{Z}/2$ coefficients, this is reminiscent of relations such as $w_2^2 \equiv p_1 \pmod{2}$. In fact, this generator is the Euler class.

Definition 3.3 *The Spin groups in the above three cases will be the pullbacks in the following diagrams:*

$$\begin{array}{ccccc}
BG(2,2) & \longrightarrow & pt & & BG(2,q) & \longrightarrow & pt & & BG(p,q) & \longrightarrow & pt \\
\downarrow \lrcorner & & \downarrow & & \downarrow \lrcorner & & \downarrow & & \downarrow \lrcorner & & \downarrow \\
BSO(2,2) & \xrightarrow{\sqrt{p_1} \times \sqrt{p_1}} & K(\mathbb{Z} \times \mathbb{Z}, 2) & , & BSO(2,q) & \xrightarrow{\sqrt{p_1} \times w_2} & K(\mathbb{Z} \times \mathbb{Z}/2, 2) & , & BSO(p,q) & \xrightarrow{w_2 \times w_2} & K(\mathbb{Z}/2 \times \mathbb{Z}/2, 2) & ,
\end{array}$$

for $p, q \geq 3$.

To justify this, we need to consider the cohomology groups. For that, we first need some calculations.

Lemma 3.4 *The following table gives the homology groups with integral coefficient $H_k(BSO(n); \mathbb{Z})$ for $k = 0, 1$ and 2 :*

	$n = 1$	$n = 2$	$n \geq 3$
H_0	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$
H_1	0	0	0
H_2	0	$\mathbb{Z}$	$\mathbb{Z}/2$

Proof. $SO(1) \cong \{1\}$ and so this case has trivial homology in nonzero degrees. Since $SO(2) \cong S^1$, we have $BSO(2) \cong CP^\infty$. We know that

$$H_k(\mathbb{C}P^n; \mathbb{Z}) = \begin{cases} \mathbb{Z}, & \text{if } 0 \leq k \leq 2n \text{ and } k \text{ is even,} \\ 0, & \text{otherwise.} \end{cases}$$

so by using the limit $H_k(\mathbb{C}P^\infty; \mathbb{Z}) = \varprojlim_n H_k(\mathbb{C}P^n; \mathbb{Z})$, we obtain

$$H_k(\mathbb{C}P^\infty; \mathbb{Z}) = \begin{cases} \mathbb{Z}, & \text{for } k \text{ even,} \\ 0, & \text{for } k \text{ odd.} \end{cases}$$

For $n \geq 3$, we know

$$\pi_k(BSO(n)) \cong \pi_{k-1}(SO(n)) = \begin{cases} 0, & \text{if } k = 1 \\ \mathbb{Z}/2, & \text{if } k = 2. \end{cases}$$

Hence, $H_k(BSO(n); \mathbb{Z})$ follows from the Hurewicz theorem. □

Proposition 3.5 *For $p, q \geq 2$, there is an isomorphism*

$$H^2(\widehat{BSO}(p, q); \mathbb{Z}) \cong H^2(BSO(p); \mathbb{Z}) \times H^2(BSO(q); \mathbb{Z}).$$

Proof. For any positive integer p and q , the Künneth formula gives

$$\begin{aligned}
H^2(BSO(p) \times BSO(q); \mathbb{Z}) &\cong \text{hom}(H_2(BSO(p) \times BSO(q); \mathbb{Z}), \mathbb{Z}) \\
&\quad \oplus \text{Ext}_{\mathbb{Z}}^1(H_1(BSO(p) \times BSO(q); \mathbb{Z}), \mathbb{Z}).
\end{aligned}$$

Now the homology groups inside the hom and Ext factors on the right hand side are calculated as

$$\begin{aligned}
H_2(BSO(p) \times BSO(q); \mathbb{Z}) &\cong \left(\bigoplus_{r+s=2} H_r(BSO(p); \mathbb{Z}) \otimes_{\mathbb{Z}} H_s(BSO(q); \mathbb{Z}) \right) \\
&\oplus \left(\bigoplus_{r+s=1} \text{Tor}_1^{\mathbb{Z}}(H_r(BSO(p); \mathbb{Z}), H_s(BSO(q); \mathbb{Z})) \right) \\
&\cong \begin{cases} 0, & \text{if } p = 1, q > 2 \text{ or } p > 2, q = 1, \\ \mathbb{Z}, & \text{if } p = 1, q = 2 \text{ or } p = 2, q = 1, \\ \mathbb{Z} \oplus \mathbb{Z} \cong H_2(BSO(p); \mathbb{Z}) \oplus H_2(BSO(p); \mathbb{Z}), & \text{if } p, q = 2 \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2 \cong H_2(BSO(p); \mathbb{Z}) \oplus H_2(BSO(p); \mathbb{Z}), & \text{if } p, q \geq 3, \end{cases}
\end{aligned}$$

and

$$\begin{aligned}
H_1(BSO(p) \times BSO(q); \mathbb{Z}) &\cong \left(\bigoplus_{r+s=1} H_r(BSO(p); \mathbb{Z}) \otimes_{\mathbb{Z}} H_s(BSO(q); \mathbb{Z}) \right) \\
&\oplus \left(\bigoplus_{r+s=0} \text{Tor}_1^{\mathbb{Z}}(H_r(BSO(p); \mathbb{Z}), H_s(BSO(q); \mathbb{Z})) \right) \\
&= 0,
\end{aligned}$$

since $H_1(BSO(n); \mathbb{Z})$ is trivial from Lemma 3.4. □

3.3 Indefinite String groups

In this section we take as our starting point the indefinite Spin group $\text{Spin}(p, q)$. We emphasize that there is a subtlety here in that this group is *not* simply connected for general p and q . In fact, the maximal compact subgroup of $\text{Spin}(p, q)$ is $\text{Spin}(p) \times \text{Spin}(q) / \{(1, 1), (-1, -1)\}$. The group $\text{Spin}(p, q)$ itself is the diagonal 2-fold cover of the 4-fold cover of $\text{SO}(p, q)$. For $p \geq q$, the fundamental group is given as

$$\pi_1(\text{Spin}(p, q)) = \begin{cases} \{0\} & (p, q) = (1, 1) \text{ or } (1, 0), \\ \{0\} & p > 2, q = 0, 1, \\ \mathbb{Z} & (p, q) = (2, 0) \text{ or } (2, 1), \\ \mathbb{Z} \times \mathbb{Z} & (p, q) = (2, 2), \\ \mathbb{Z} & p > 2, q = 2, \\ \mathbb{Z}/2 & p, q > 2. \end{cases} \quad (3.9)$$

In order to define indefinite String structure properly we need to take as a starting point a simply-connected group. Therefore, we should start from the simply-connected cover of $\text{Spin}(p, q)$, which is what we do below. Note, however, that one can define variants of String structures without requiring this. The resulting structure would be analogous to the case of p_1 -structures (see [24] [25] for various analogous structures in the definite case).

We next kill the next nontrivial homotopy groups, namely π_3 , of the relevant Spin group $\text{Spin}(p, q)$. Note that at this stage if either p or q is less than 3 then the corresponding factor

in the decomposition $\text{Spin}(p) \times \text{Spin}(q)$ will not be seen in the process. So we will consider mainly two cases: $\text{Spin}(n) \simeq \widehat{\text{Spin}}(1, n) \simeq \widehat{\text{Spin}}(2, n) \simeq \text{O}(n)\langle 3 \rangle$ and $\widehat{\text{Spin}}(p, q) \simeq \text{O}(p, q)\langle 3 \rangle$ for $p, q \geq 3$.

When $p, q \geq 3$, the maximal compact subgroup of the 1-connected cover $\text{O}(p, q)\langle 1 \rangle$ is $\text{O}(p)\langle 1 \rangle \times \text{O}(q)\langle 1 \rangle$ which is homotopy equivalent to $\text{Spin}(p) \times \text{Spin}(q)$. Hence, $\text{O}(p, q)\langle 1 \rangle$ is homotopy equivalent to $\text{Spin}(p) \times \text{Spin}(q)$. In fact, since $\text{O}(1)\langle 1 \rangle$ and $\text{O}(2)\langle 1 \rangle$ are just a point space and contractible space respectively, we still can say that $\text{O}(p, q)\langle 1 \rangle$ is homotopy equivalent to $\text{O}(p)\langle 1 \rangle \times \text{O}(q)\langle 1 \rangle$.

First we will show that $H^4(B\text{Spin}(n); \mathbb{Z}) \cong \mathbb{Z}$ with a unique generator $\frac{1}{2}p_1$ such that $\frac{1}{2}p_1 + \frac{1}{2}p_1 = p_1$ is the first Pontrjagin class. Although this has been used in many papers, and we are certain that it is known to experts, we could not find an explicit written proof of this in the literature. The results in this direction are complicated and highly dependent on n – the reader might wish to consult the papers [5] [8] [15] [3] to appreciate the subtleties involved. We will use these cited works to deduce the result. Note that, in contrast, the fact that $\frac{1}{2}p_1$ in the stable case generates $H^4(B\text{Spin}; \mathbb{Z})$ follows directly from the general result on the ring structure [32].

By [15], the torsion elements of $H^*(B\text{Spin}(n); \mathbb{Z})$ are of order 2. Furthermore, as an abelian group

$$H^k(B\text{Spin}(n); \mathbb{Z}) = \mathbb{Z}^{q_{n,k}} \oplus (\mathbb{Z}/2)^{r_{n,k}}, \quad (3.10)$$

where $q_{n,k}$ is the k th Betti number and $r_{n,k}$ are analogously defined numbers (the explicit expression of which we do not need). Expanding the generating function $Q_n(t) := \sum_{k=0}^{\infty} q_{n,k} t^k \in \mathbb{Z}[[t]]$ defined by

$$Q_{2n}(t) = \left(\prod_{j=1}^{n-1} (1 - t^{4j}) \right)^{-1} (1 - t^{2n})^{-1}, \quad Q_{2n+1}(t) = \left(\prod_{j=1}^n (1 - t^{4j}) \right)^{-1}, \quad (3.11)$$

we have that $q_{n,4} = 1$. Similarly, it can be shown that $r_{n,k} = 0$.

Let $\rho : T \hookrightarrow \text{Spin}(n)$ be the embedding of a maximal torus, and $\rho_* : BT \rightarrow B\text{Spin}(n)$ be the corresponding map of classifying spaces. The induced map on integral cohomology is the restriction map $\rho^* : H^*(B\text{Spin}(n); \mathbb{Z}) \rightarrow H^*(BT; \mathbb{Z})$, whose kernel is the ideal of torsion elements in $H^*(B\text{Spin}(n); \mathbb{Z})$ [4]. The images lies in the Weyl invariants and so ρ^* gives rise to a well-defined injective map

$$\bar{\rho}^* : H^*(B\text{Spin}(n); \mathbb{Z})/\text{torsion} \longrightarrow H^*(BT; \mathbb{Z})^W, \quad (3.12)$$

where the range is the invariants of the Weyl group of the integral cohomology ring of the classifying space of the maximal torus. While $\bar{\rho}^*$ is an isomorphism for $BSO(n)$ [5] [8] this is not the case for $B\text{Spin}(n)$ [3]. Furthermore, as for the ring structure, $H^*(B\text{Spin}(n); \mathbb{Z})$ is isomorphic to the subring of

$$\ker \beta \oplus H^*(B\text{Spin}(n); \mathbb{Z})/\text{torsion} \quad (3.13)$$

consisting of pairs of elements with the same image in $\mathbb{Z}/2 \otimes_{\mathbb{Z}} (H^*(B\text{Spin}(n); \mathbb{Z})/\text{torsion})$ [3]. Here $\beta = Sq^1$ is the mod 2 Bockstein operation.

We now identify the generator, putting together various results in [3]. One might in principle have classical classes as well as new exotic classes. First, the Pontrjagin classes $p_i \in H^*(BT; \mathbb{Z})^W$ are in the images of $\bar{\rho}^*$ because they are just obtained from $BSO(n)$. The lowest degree, which is 4, resembles to a large extent the stable case, namely there are elements $q_i \in H^*(BT; \mathbb{Z})^W$ ($i \geq 1$) such that $\deg(q_i) = 2^{i+1}$ with a relation in lowest degree given by $2q_1 = p_1$ (Proposition 3.3 in [3]). The cohomology ring is given by $H^*(BT; \mathbb{Z})^W = \mathbb{Z}[p_1, q_1, \dots]$ with identified generators, and with

p_1 being redundant since $p_1 = 2q_1$ (see proof of Corollary 7.2 in [3]). Similarly, the Euler class e which appears for even n is also redundant as e is proportional to p_1 in degree four. Therefore, q_1 is the only generator of $H^4(BT; \mathbb{Z})^W$.

The kernel of Sq^1 on $H^*(B\text{Spin}(n); \mathbb{Z}/2)$ is the reduction mod 2 of $H^*(B\text{Spin}(n); \mathbb{Z})$ and its image is the reduction mod 2 of the torsion elements of $H^*(B\text{Spin}(n); \mathbb{Z})$ [15]. Now the element $p_1 \in H^*(BSO(n); \mathbb{Z})$ reduces modulo 2 to the square $w_2 \cup w_2 \in H^*(BSO(n); \mathbb{Z}/2)$, which maps to zero in $H^*(B\text{Spin}(n); \mathbb{Z}/2)$. This means that p_1 is divisible by 2 in $H^*(B\text{Spin}(n); \mathbb{Z}/2)$, i.e. $p_1 = 2q'_1$ for some q'_1 . Then we have $\bar{p}^*(2q'_1) = 2q_1$ (see [3]). Now because $H^*(BT; \mathbb{Z})^W$ is torsion-free we then have $\bar{p}^*(q'_1) = q_1$. Since q_1 is not divisible by 2 then neither is q'_1 . This implies that the reduction modulo 2 of q'_1 is nonzero.

A further argument is that, from [15], an element of $H^*(B\text{Spin}(n); \mathbb{Z})$ is completely determined by its canonical images in $H^*(B\text{Spin}(n); \mathbb{R})$ and $H^*(B\text{Spin}(n); \mathbb{Z}/2)$. Now, from [22], the structure of mod 2 cohomology of the unstable Spin group is given by

$$\begin{aligned} H^*(B\text{Spin}(n); \mathbb{Z}/2) &= H^*(BSO(n); \mathbb{Z}/2)/J_h \otimes \mathbb{Z}/2[w_{2^h}(\Delta)] \\ &= \mathbb{Z}/2[w_2, \dots, w_n]/(r_1, \dots, r_h) \otimes \mathbb{Z}/2[w_{2^h}(\Delta)] , \end{aligned}$$

where Δ is a real Spin representation of $\text{Spin}(n)$ and h denotes certain integers related to the dimension (mod 2) of the representation. The elements $r_j \in H^{2^{j-1}+1}(BSO(n); \mathbb{Z}/2)$ appearing in the module J_h are defined by $r_1 = w_2$ and $r_{j+1} = Sq^{2^{j-1}}r_j$ for $j \geq 1$. We are interested in cohomology degree four, and so the relevant pieces of data will be $r_1 = w_2$, $r_2 = Sq^1w_2 = w_3$ and $r_3 = Sq^2w_2$, while the Spin representation term will not be relevant by dimension reason, as for us $h > 2$. The truncated polynomial ring is $\mathbb{Z}/2[w_2, w_3, w_4, \dots]$, while the module is given by $(w_2, w_3, Sq^2w_2, \dots)$, which implies that overall we have that

$$H^4(B\text{Spin}(n); \mathbb{Z}/2) \cong \mathbb{Z}/2[w_4] . \quad (3.14)$$

Finally, note that w_4 is the mod 2 reduction of $\frac{1}{2}p_1$.

Therefore, we have

Proposition 3.6 *The fourth cohomology group is $H^4(B\text{Spin}(n); \mathbb{Z}) \cong \mathbb{Z}$ and is generated by $\frac{1}{2}p_1$.*

We now get back to the construction of indefinite String structures. We obtain a 3-connected cover $G(n)$ for $n \geq 3$ by taking pullbacks, contrasting two cases:

$$\begin{array}{ccc} BG(n) & \xrightarrow{\quad} & \text{pt} \\ \downarrow \lrcorner & & \downarrow \\ B\text{Spin}(n \neq 4) & \xrightarrow{\frac{1}{2}p_1} & K(\mathbb{Z}, 4) \end{array} \quad \begin{array}{ccc} BG(4) & \xrightarrow{\quad} & \text{pt} \\ \downarrow \lrcorner & & \downarrow \\ B\text{Spin}(4) & \xrightarrow{(\frac{1}{2}p_1, \frac{1}{2}p_1)} & K(\mathbb{Z} \times \mathbb{Z}, 4) . \end{array} \quad (3.15)$$

Here, the map $(\frac{1}{2}p_1, \frac{1}{2}p_1) : B\text{Spin}(4) \rightarrow K(\mathbb{Z} \times \mathbb{Z}, 4)$ is equivalent to the map $\frac{1}{2}p_1 \times \frac{1}{2}p_1 : B\text{Spin}(3) \times B\text{Spin}(3) \rightarrow K(\mathbb{Z} \times \mathbb{Z}, 4)$ through the accidental isomorphism $\text{Spin}(4) \cong \text{Spin}(3) \times \text{Spin}(3)$. Consequently, any classifying map $\widehat{^2 f} : X \rightarrow \widehat{B\text{Spin}(3, 3)}$ can be decomposed into a pair $f = (f_1, f_2)$ with $f_1, f_2 : B\text{Spin}(3) \rightarrow K(\mathbb{Z}, 4)$. Recall also that for any classifying map $f : X \rightarrow \widehat{B\text{Spin}(p, q)}$

²Here a hat on the group indicates that we are taking the simply connected cover.

can be decomposed into $(f_1, f_2) : X \rightarrow B\widehat{\text{Spin}}(p) \times B\widehat{\text{Spin}}(q)$. Hence we can use the additive and multiplicative notations interchangeably.

In order to kill π_3 of $\widehat{\text{Spin}}(p, q)$, we take the pullbacks according to the following procedure.

Definition 3.7 *The String groups in the indefinite case are defined via the following pullbacks:*
(i) for $p, q \geq 5$

$$\begin{array}{ccc} B\widehat{\text{String}}(p, q) & \xrightarrow{\quad} & \text{pt} \\ \downarrow & & \downarrow \\ B\widehat{\text{Spin}}(p, q) & \xrightarrow{\frac{1}{2}p_1 \times \frac{1}{2}p_1} & K(\mathbb{Z} \times \mathbb{Z}, 4) ; \end{array} \quad (3.16)$$

(ii) for $p = 4, q \geq 5$,

$$\begin{array}{ccc} B\widehat{\text{String}}(4, q) & \xrightarrow{\quad} & \text{pt} \\ \downarrow & & \downarrow \\ B\widehat{\text{Spin}}(4, q) & \xrightarrow{(\frac{1}{2}p_1 \times \frac{1}{2}p_1) \times \frac{1}{2}p_1} & K(\mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}, 4) ; \end{array} \quad (3.17)$$

(iii) and for $p = q = 4$,

$$\begin{array}{ccc} B\widehat{\text{String}}(4, 4) & \xrightarrow{\quad} & \text{pt} \\ \downarrow & & \downarrow \\ B\widehat{\text{Spin}}(4, 4) & \xrightarrow{(\frac{1}{2}p_1 \times \frac{1}{2}p_1) \times (\frac{1}{2}p_1 \times \frac{1}{2}p_1)} & K(\mathbb{Z} \times \mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}, 4) . \end{array} \quad (3.18)$$

Note that we can also consider variants of String structures associated to the non-simply connected groups $\text{Spin}(p, q)$.

Definition 3.8 *A (p_1, p'_1) -structure is a lift from $B\text{Spin}(p, q)$ to the classifying space obtained by killing the fourth homotopy group.*

These are analogs of a p_1 -structure, where the lower homotopy groups are not necessarily killed. See [24] [25] [26] for extensions and applications.

Of course, we need to establish a decomposition of the corresponding cohomology groups. This in turn will require taking homomorphisms with homology.

Remark 5 *We will see shortly (in the proof of Proposition 3.11 below) that $H^4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}) \cong H^4(B\text{Spin}(p); \mathbb{Z}) \oplus H^4(B\text{Spin}(q); \mathbb{Z})$. One might be tempted to use the following relation, for an arbitrary space X and Y and any k ,*

$$H^k(X \times Y; R) \cong \bigoplus_{r+s=k} H^r(X; R) \otimes_R H^s(Y; R)$$

which holds when R is a field. Unfortunately, we can not do this in our case since $\mathbb{Z}$ is not a field. However, we may use the Künneth formula and the universal coefficient theorem because $\mathbb{Z}$ is a principal ideal domain. This involves the computation of homology groups and related Tor and Ext groups. While these are straightforward, we find it unavoidable to go through them to ensure that we are on firm ground in this surprisingly delicate setting of indefinite structures.

To that end, we start with the following:

Lemma 3.9

$$H_4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}) \cong \begin{cases} H_4(B\text{Spin}(q); \mathbb{Z}) & \text{for any } q \text{ if } p = 1, \\ H_4(B\text{Spin}(p); \mathbb{Z}) & \text{for any } p \text{ if } q = 1, \\ H_4(B\text{Spin}(p); \mathbb{Z}) \oplus H_4(B\text{Spin}(q); \mathbb{Z}) & \text{if } p, q \geq 2. \end{cases}$$

Proof. The Künneth formula for homology gives the following identity:

$$\begin{aligned} H_4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}) \cong & \left(\bigoplus_{r+s=4} H_r(B\text{Spin}(p); \mathbb{Z}) \otimes_{\mathbb{Z}} H_s(B\text{Spin}(q); \mathbb{Z}) \right) \\ & \oplus \left(\bigoplus_{r+s=3} \text{Tor}_1^{\mathbb{Z}}(H_r(B\text{Spin}(p); \mathbb{Z}), H_s(B\text{Spin}(q); \mathbb{Z})) \right). \end{aligned}$$

Since $H_s(B\text{Spin}(q); \mathbb{Z}) = 0$ for $s = 1, 2, 3$, the only nontrivial term in Tor is

$$\text{Tor}_1^{\mathbb{Z}}(H_3(B\text{Spin}(p); \mathbb{Z}), H_0(B\text{Spin}(q); \mathbb{Z})).$$

This is also trivial for $p \geq 2$. Moreover, when $p = 1$, we have $\text{Tor}_1^{\mathbb{Z}}(\mathbb{Z}/2, \mathbb{Z})$ and this is trivial since $\mathbb{Z}$ is torsion-free.

The direct sum term on the right hand side of the above Künneth formula has only two nontrivial factors: $H_0(B\text{Spin}(p); \mathbb{Z}) \otimes H_4(B\text{Spin}(q); \mathbb{Z})$ and $H_4(B\text{Spin}(p); \mathbb{Z}) \otimes H_0(B\text{Spin}(q); \mathbb{Z})$. When $p \geq 2$, these two are isomorphic to $\mathbb{Z} \oplus \mathbb{Z} \cong \mathbb{Z}$. At this stage, there seems to be several routes to take. Since we do not want to lose the structure of homology group of degree four, we *choose* the isomorphisms

$$\begin{aligned} H_0(B\text{Spin}(p); \mathbb{Z}) \otimes H_4(B\text{Spin}(q); \mathbb{Z}) &\cong H_4(B\text{Spin}(q); \mathbb{Z}), \\ H_4(B\text{Spin}(p); \mathbb{Z}) \otimes H_0(B\text{Spin}(q); \mathbb{Z}) &\cong H_4(B\text{Spin}(p); \mathbb{Z}). \end{aligned}$$

On the other hand, when $p = 1$, we have $H_4(B\text{Spin}(1); \mathbb{Z}) = 0$. So the only nontrivial term is now $H_0(B\text{Spin}(p); \mathbb{Z}) \otimes H_4(B\text{Spin}(q); \mathbb{Z}) \cong H_4(B\text{Spin}(q); \mathbb{Z})$. $\square$

We will also need to calculate the Ext-term.

Lemma 3.10

$$\text{Ext}_{\mathbb{Z}}^1(H_3(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}), \mathbb{Z}) \cong \begin{cases} \mathbb{Z}/2 & \text{if } p = 1 \text{ or } q = 1, \\ 0 & \text{if } p, q \geq 2. \end{cases}$$

Proof. First, we need to compute $H_3(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z})$ and we use the Künneth formula:

$$\begin{aligned} H_3(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}) \cong & \left(\bigoplus_{r+s=3} H_r(B\text{Spin}(p); \mathbb{Z}) \otimes_{\mathbb{Z}} H_s(B\text{Spin}(q); \mathbb{Z}) \right) \\ & \oplus \left(\bigoplus_{r+s=2} \text{Tor}_1^{\mathbb{Z}}(H_r(B\text{Spin}(p); \mathbb{Z}), H_s(B\text{Spin}(q); \mathbb{Z})) \right). \end{aligned}$$

The Tor term is trivial since $H_s(B\text{Spin}(q); \mathbb{Z}) = 0$ or $\mathbb{Z}$, and $\mathbb{Z}$ is torsion-free. The only nontrivial factor in the first term on the right hand side is $H_3(B\text{Spin}(p); \mathbb{Z}) \otimes H_0(B\text{Spin}(q); \mathbb{Z})$. This is zero for $p \geq 2$ since $H_3(B\text{Spin}(p); \mathbb{Z}) = 0$. On the other hand, if $p = 1$, we have $H_3(B\text{Spin}(1); \mathbb{Z}) = \mathbb{Z}/2$ and the result follows from the relation $\mathbb{Z}/2 \otimes \mathbb{Z} \cong \mathbb{Z}/2$. $\square$

We are now ready to calculate the degree four cohomology groups.

Proposition 3.11 *For any $p, q \geq 2$, we have*

$$H^4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}) \cong H^4(B\text{Spin}(p); \mathbb{Z}) \oplus H^4(B\text{Spin}(q); \mathbb{Z}) .$$

For the lower dimensional case, we have

$$H^4(B\text{Spin}(1) \times B\text{Spin}(q); \mathbb{Z}) \cong H^4(B\text{Spin}(q); \mathbb{Z}) .$$

Proof. The Künneth formula for cohomology asserts that

$$\begin{aligned} H^4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}) &\cong \text{hom}(H_4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}), \mathbb{Z}) \\ &\quad \oplus \text{Ext}_{\mathbb{Z}}^1(H_3(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}), \mathbb{Z}) . \end{aligned}$$

When $p = 1$, by the above lemmas, we have

$$\begin{aligned} H^4(B\text{Spin}(1) \times B\text{Spin}(q); \mathbb{Z}) &\cong \text{hom}(H_4(B\text{Spin}(q); \mathbb{Z}) \oplus \mathbb{Z}/2, \mathbb{Z}) \\ &\cong H^4(B\text{Spin}(1); \mathbb{Z}) \times H^4(B\text{Spin}(q); \mathbb{Z}) . \end{aligned}$$

Here, we used the fact that finite products and finite coproducts coincide in the additive category. When $p \geq 2$, we have

$$\begin{aligned} H^4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z}) &\cong \text{hom}(H_4(B\text{Spin}(p); \mathbb{Z}) \oplus H_4(B\text{Spin}(q); \mathbb{Z}), \mathbb{Z}) \\ &\cong \text{hom}(H_4(B\text{Spin}(p); \mathbb{Z}), \mathbb{Z}) \times \text{hom}(H_4(B\text{Spin}(q); \mathbb{Z}), \mathbb{Z}) \\ &\cong H^4(B\text{Spin}(p); \mathbb{Z}) \times H^4(B\text{Spin}(q); \mathbb{Z}) . \end{aligned}$$

For $p = 2$, we have $H_4(B\text{Spin}(2); \mathbb{Z}) \cong H_4(\mathbb{CP}^\infty; \mathbb{Z}) \cong \mathbb{Z}$, so that $H^4(B\text{Spin}(2); \mathbb{Z}) \cong \mathbb{Z}$.

The maximal compact subgroup of $G = \text{Spin}(p, q)$ is $K = \text{Spin}(p) \times \text{Spin}(q)/(\mathbb{Z}/2)$. Therefore, G and K are weakly homotopy equivalent to each other so that they are in fact homotopy equivalent, since the usual cohomology is represented by the Eilenberg-MacLane spaces in a sense that $H^n(X; A) \cong [X, K(A, n)]$, where A is a coefficient group (or an integer ring) and X is arbitrary topological space. From the following short exact sequence

$$1 \rightarrow \mathbb{Z}/2 \rightarrow \text{Spin}(p) \times \text{Spin}(q) \rightarrow (\text{Spin}(p) \times \text{Spin}(q))/(\mathbb{Z}/2) \rightarrow 1$$

and the fact that $H^n(\mathbb{Z}/2; \mathbb{Z}) = 0$ for any n , we obtain the desired isomorphism $H^4(B\text{Spin}(p, q); \mathbb{Z}) \cong H^4(B\text{Spin}(p) \times B\text{Spin}(q); \mathbb{Z})$. $\square$

3.4 String structure associated to indefinite unitary and symplectic groups

The indefinite unitary group. Let $U(p, q)$ denote the group of matrices of linear isometries of the pseudo-Hermitian space $\mathbb{C}^{p, q}$ of signature p, q . The special indefinite unitary group $SU(p, q) = U(p, q) \cap \text{SL}(p + q, \mathbb{C})$ is the subgroup of $U(p, q)$ consisting of matrices of determinant 1.

The following table lists the homotopy groups of the unitary group [13] (see [18] [21] for explicit generators).

	U(1)	U(2)	U(3)	U(4)	U(5)	U(6)
π_1	$\boxed{\mathbb{Z}}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$
π_2	0	0	0	0	0	0
π_3	0	$\boxed{\mathbb{Z}}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$
π_4	0	$\mathbb{Z}/2$	0	0	0	0
π_5	0	$\mathbb{Z}/2$	$\boxed{\mathbb{Z}}$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$
π_6	0	$\mathbb{Z}/12$	$\mathbb{Z}/6$	0	0	0
π_7	0	$\mathbb{Z}/2$	0	$\boxed{\mathbb{Z}}$	$\mathbb{Z}$	$\mathbb{Z}$

Remark 6 (i) *The indefinite unitary group admits a Cartan decomposition $U(p, q) \cong U(p) \times U(q) \times \mathbb{C}^{pq}$, so that – as in the case of the orthogonal group – the cohomology is determined by the maximal compact subgroup $K = U(p) \times U(q)$. This gives rise to two cohomology classes, one from each factor in K , except when p or q is equal to 1, in which case there is only one class in real degree four, namely c_2 for the complementary nontrivial factor in $U(1, q)$ or $U(p, 1)$.*

(ii) *All the unitary groups have nontrivial fundamental group, isomorphic to $\mathbb{Z}$. The universal covering groups of the indefinite unitary group and the special indefinite unitary group are denoted $\widetilde{U}(p, q)$ and $\widetilde{SU}(p, q)$, respectively. Note that the latter is also a subgroup of the former.*

(iii) *As far as π_3 is concerned, the groups are already in the stable range. This makes the discussion much simpler than in the orthogonal case.*

(iv) *The cohomology rings of classifying spaces of the unitary group and the special unitary group with integral coefficients are generated by the Chern classes c_i in degree $2i$ and has a considerably simpler form than the orthogonal case, i.e.*

$$\begin{aligned} H^*(BU(n); \mathbb{Z}) &\cong \mathbb{Z}[c_1, c_2, c_3, \dots, c_n], \\ H^*(BSU(n); \mathbb{Z}) &\cong \mathbb{Z}[c_2, c_3, \dots, c_n]. \end{aligned}$$

String structures associated to the unitary group are considered in constructions associated to elliptic cohomology, such as in [1].

Definition 3.12 *A String structure for the indefinite unitary group $U(p, q)$ is defined via the following pullback diagrams*

(i) *for $p, q \geq 2$*

$$\begin{array}{ccc} \widehat{BString}(U(p, q)) & \longrightarrow & \text{pt} \\ \downarrow \lrcorner & & \downarrow \\ \widehat{BU}(p, q) & \xrightarrow{c_2 \times c'_2} & K(\mathbb{Z} \times \mathbb{Z}, 4); \end{array} \quad (3.19)$$

(ii) *for $p = 1$*

$$\begin{array}{ccc} \widehat{BString}(U(1, q)) & \longrightarrow & \text{pt} \\ \downarrow \lrcorner & & \downarrow \\ \widehat{BU}(1, q) & \xrightarrow{c'_2} & K(\mathbb{Z}, 4); \end{array} \quad (3.20)$$

and similarly for $q = 1$ with a class c_2 corresponding to the first factor.

The following is immediate from the definition.

Proposition 3.13 (i) $\text{String}(\text{U}(p, q))$ and $\text{String}(\text{SU}(p, q))$ structures are classified by a pair of classes (c_2, c'_2) , where c_2 and c'_2 are the generators in degree 4 of $\text{BU}(p)$ and $\text{BU}(q)$, respectively.
(ii) When either $p = 1$ or $q = 1$, we only have one generator as an obstruction for $\text{String}(\text{U}(p, q))$.

The indefinite symplectic group. The indefinite symplectic group $\text{Sp}(p, q)$, also known as the indefinite quaternionic unitary group $\text{U}(p, q; \mathbb{H})$, can be defined as the isometry group of a nondegenerate quaternionic Hermitian form in $\mathbb{H}^n$.

Remark 7 (i) The indefinite symplectic group admits a Cartan decomposition $\text{Sp}(p, q) \cong \text{Sp}(p) \times \text{Sp}(q) \times \mathbb{H}^{pq}$, so that again the cohomology is determined by the maximal compact subgroup $K = \text{Sp}(p) \times \text{Sp}(q)$, giving rise to two cohomology classes. Furthermore, due to the relatively large dimension, there are no degenerate cases here. For instance, $\text{Sp}(1, 1) \cong \text{Sp}(1) \times \text{Sp}(1) \cong S^3 \times S^3$.

(ii) The symplectic group is simply connected, so there are no issues with the starting point to define a corresponding String structure.

(iii) $\pi_i(\text{Sp}(n))$ is in the stable range already for $i \leq 4n + 1$. Therefore, we are in the stable range for any value of n when considering the third homotopy group. The homotopy groups of the symplectic groups are computed by Mimura and Toda (see [19]).

(iv) The cohomology ring of the classifying space of the symplectic group is generated by the symplectic Pontrjagin classes $p_i^{\mathbb{H}}$ of degree $4i$,

$$H^*(B\text{Sp}(n); \mathbb{Z}) \cong \mathbb{Z}[p_1^{\mathbb{H}}, p_2^{\mathbb{H}}, \dots, p_n^{\mathbb{H}}]. \quad (3.21)$$

(v) Under the identification $\text{Sp}(1) \cong \text{SU}(2)$, $p_1^{\mathbb{H}}$ is equal to $-c_2$.

Definition 3.14 A String structure for the indefinite symplectic group $\text{Sp}(p, q)$ is defined via the pullback

$$\begin{array}{ccc} B\text{String}(\text{Sp}(p, q)) & \xrightarrow{\quad} & \text{pt} \\ \downarrow & & \downarrow \\ B\text{Sp}(p, q) & \xrightarrow{p_1^{\mathbb{H}} \times p_1^{\mathbb{H}}} & K(\mathbb{Z} \times \mathbb{Z}, 4) \end{array} \quad (3.22)$$

As in the unitary case, it follows directly from the definition that we have the following.

Proposition 3.15 String structures associated with the indefinite symplectic group are classified by a pair of symplectic Pontrjagin classes $(p_1^{\mathbb{H}}, p_1^{\mathbb{H}'})$, where the first and second are generators of $H^4(B\text{Sp}(p); \mathbb{Z})$ and $H^4(B\text{Sp}(q); \mathbb{Z})$, respectively.

3.5 Relation to twisted structures

We have seen that indefinite structures are determined homotopically by their maximal compact subgroups that are products of two compact Lie groups. The obstruction encountered involves two characteristic classes, one from each of these factor group. It is then natural to investigate how the two ‘composite structures’ might interact. There is another instance where a pair of cohomological

structures interact in this context, namely twisted structures ([37] [28] [24] [25] [26]), to which we now explore possible connections.

We have seen that a G -principal bundle $f : X \rightarrow BG$ with a topological group G that is homotopy equivalent to $G' \times G''$ can be lifted to a $\tilde{G}$ -principal bundle, where $\tilde{G}$ has the same homotopy type as that of G except that π_n killed, when both obstruction classes $f_1^* \alpha'$ and $f_2^* \alpha''$ in $H^{n+1}(X; \pi_n(G' \times G''))$ vanish. That is, the outer square in the following diagram commutes up to homotopy:

$$\begin{array}{ccc}
 X & \xrightarrow{\quad} & \text{pt} \\
 \searrow^{f=(f_1, f_2)} & \nearrow & \downarrow \\
 & B\tilde{G} & \\
 & \downarrow & \\
 B(G' \times G'') & \xrightarrow{\alpha' \times \alpha''} & K(\pi_n(G' \times G''_{n-1}), n+1) .
 \end{array} \tag{3.23}$$

Instead of requiring both obstructions $f_1^* \alpha'$ and $f_2^* \alpha''$ to vanish simultaneously, we may want to relax this condition to just the vanishing of the difference

$$f_1^* \alpha' - f_2^* \alpha'' = 0 \in H^{n+1}(X; \pi_n(G' \times G'')). \tag{3.24}$$

Now suppose the two groups G' and G'' have the same homotopy groups in degree n , i.e. we have $\pi_n(G') \cong \pi_n(G'')$. Denoting this isomorphism group by π_n , we take the pullback $\tilde{G}_0$ as the following diagram:

$$\begin{array}{ccc}
 \tilde{G}_0 & \xrightarrow{\quad} & BG'' \\
 \downarrow \lrcorner & & \downarrow \alpha'' \\
 BG' & \xrightarrow{\alpha'} & K(\pi_n, n+1) .
 \end{array} \tag{3.25}$$

Suppose that $\tilde{G} := \Omega \tilde{G}_0$ has a topological group structure. Given $f_1 : X \rightarrow BG'$ and $f_2 : X \rightarrow BG''$ classifying G' - and G'' -principal bundles over X , respectively, there is a universal $\tilde{G}$ -principal bundle over X if $f_1^* \alpha'$ is homotopic to $f_2^* \alpha''$ which is equivalent to the condition

$$f_1^* \alpha' - f_2^* \alpha'' = 0 \in H^{n+1}(X; \pi_n) . \tag{3.26}$$

Diagrammatically, the condition is equivalent to requiring that there be a homotopy h as indicated in the diagram:

$$\begin{array}{ccc}
 X & \xrightarrow{f_2} & BG'' \\
 \searrow^{f_1} & \nearrow & \downarrow \alpha'' \\
 & B\tilde{G} & \\
 & \downarrow & \\
 BG' & \xrightarrow{\alpha'} & K(\pi_n, n+1) .
 \end{array} \tag{3.27}$$

(Note: In the original image, a homotopy h is indicated between the maps $B\tilde{G} \rightarrow BG''$ and $B\tilde{G} \rightarrow BG'$.)

All of this motivates the following definition:

Definition 3.16 Suppose that we have two topological groups G' and G'' with homotopy groups $\pi'_n := \pi_n(G')$ and $\pi''_n := \pi_n(G'')$ respectively. Moreover, suppose that we have given cohomology classes $\alpha' \in H^{n+1}(BG'; \pi'_n)$ and $\alpha'' \in H^{n+1}(BG''; \pi''_n)$ and a group homomorphism $\varphi : \pi''_n \rightarrow \pi'_n$. Then we have the homotopy group $\tilde{G}$ as in the previous argument, and for two G' - and G'' -structures over X given by f_1 and f_2 , the induced $\tilde{G}$ -structure over X is said to be twisted in favor of G' :

$$\begin{array}{ccc}
 X & \xrightarrow{f_2} & BG'' \\
 \searrow f_1 & \swarrow & \downarrow \varphi \circ \alpha'' \\
 & B\tilde{G} & \longrightarrow BG'' \\
 & \downarrow & \\
 & BG' & \xrightarrow{\alpha'} K(\pi_n, n+1) .
 \end{array} \tag{3.28}$$

Remark 8 The twisted construction has natural motivations arising from physics, as presented by Sati-Schreiber-Stasheff [28]. For instance, the Green-Schwarz anomaly condition ([10] [9])

$$\frac{1}{2}p_1(TX) - \text{ch}_2(E) = 0 \in H^4(X; \mathbb{Z}) , \tag{3.29}$$

where $\text{ch}_2(E)$ is the second Chern character of a bundle E which reduces to the second Chern class $c_2(E)$ is equivalent to existence of a homotopy H_3 in the following diagram, with $\pi_3(\text{SU}(n)) \cong \pi_3(\text{Spin}(n)) \cong \mathbb{Z}$ for $n \geq 3$ except 4,

$$\begin{array}{ccc}
 X & \xrightarrow{E} & \text{BSU}(n) \\
 \downarrow TX & \swarrow H_3 & \downarrow c_2 \\
 \text{BSpin}(n) & \xrightarrow{\frac{1}{2}p_1} & K(\mathbb{Z}, 4) .
 \end{array} \tag{3.30}$$

The homotopy H_3 exhibits the B-field as a twisted gerbe, whose twist is the difference class $\frac{1}{2}p_1(TX) - c_2(E)$. Our definition above extends this to the indefinite signature case. What we have in our current context is what might essentially be viewed as a twisted String structure, in the sense of [37] [28], where the twist itself arises from a Spin bundle, where the two are the two parts in the composite maximal compact subgroup of $\text{Spin}(p, q)$.

Remark 9 We have only two cases to consider for the twisted coverings:

$$\begin{array}{ccc}
 \widetilde{\text{BSpin}}(2, 2) & \longrightarrow & \text{BSO}(2) \\
 \downarrow & & \downarrow \sqrt{p_1} \\
 \text{BSO}(2) & \xrightarrow{\sqrt{p_1}} & K(\mathbb{Z}, 2)
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 \widetilde{\text{BSpin}}(p, q) & \longrightarrow & \text{BSO}(q) \\
 \downarrow & & \downarrow w_2 \\
 \text{BSO}(p) & \xrightarrow{w_2} & K(\mathbb{Z}/2, 2)
 \end{array}
 \quad \text{for } p, q \geq 3.$$

Similarly to the previous cases, we can construct twisted coverings:

Definition 3.17 The twisted covering indefinite String groups are defined as

$$\begin{array}{ccc}
 \widehat{\text{BString}}(p, q) & \longrightarrow & \text{BString}(q) \\
 \downarrow & & \downarrow \frac{1}{2}p_1 \\
 \text{BString}(p) & \xrightarrow{\frac{1}{2}p_1} & K(\mathbb{Z}, 4)
 \end{array}
 \quad , \quad
 \begin{array}{ccc}
 \widehat{\text{BString}}(4, 4) & \longrightarrow & \text{BString}(4) \\
 \downarrow & & \downarrow \frac{1}{2}p_1 \times \frac{1}{2}p_1 \\
 \text{BString}(4) & \xrightarrow{\frac{1}{2}p_1 \times \frac{1}{2}p_1} & K(\mathbb{Z} \times \mathbb{Z}, 4) .
 \end{array}$$

It would be very interesting to extend the definitions and constructions that we presented in this paper for String structures to include ‘pseudo-Riemannian versions’ of Fivebrane [27] [28] and Ninebrane structures [26]. This might require considerable calculations. We believe that it would also be worthwhile to explore geometric applications to gerbes, loop spaces, parallel transport, Chern-Simons theories, and stacky constructions, just to name a few. We hope to explore these topics elsewhere. Our initial goal was to get to these topics directly. However, we realized that seemingly straightforward matters are in fact much more subtle than meets the eye, so we believe it is worth addressing those first in this paper to provide firm ground from which to pursue further constructions.

Acknowledgements

We would like to thank Domenico Fiorenza, Corbett Redden, Jonathan Rosenberg, and Urs Schreiber for very useful discussions. The research of H. S. was supported by NSF Grant PHY-1102218.

References

- [1] M. Ando, M. J. Hopkins, and N. P. Strickland, *Elliptic spectra, the Witten genus and the theorem of the cube*, Invent. Math. **146** (2001), no. 3, 595–687.
- [2] H. Baum, *Spin-Strukturen und Dirac-Operatoren über pseudoriemannschen Mannigfaltigkeiten*, B.G. Teubner. Leipzig, 1981.
- [3] D. J. Benson and J. A. Wood, *Integral invariants and cohomology of $B\mathrm{Spin}(n)$* , Topology **34**, no. 1, (1995), 13–28.
- [4] A. Borel, *Topics in the homology theory of fibre bundles*, Lecture Notes in Mathematics **26**, Springer-Verlag (1967).
- [5] E. H. Brown, Jr., *The cohomology of BSO_n and BO_n with integer coefficients*, Proc. Amer. Math. Soc. **85**, no. 2, (1982), 283–288.
- [6] U. Bunke, *String structures and trivialisations of a Pfaffian line bundle*, [arXiv:0909.0846] [math.KT].
- [7] D. Fiorenza, H. Sati, and U. Schreiber, *Multiple $M5$ -branes, String 2-connections, and 7d non-abelian Chern-Simons theory*, Adv. Theor. Math. Phys. **18** (2014), 1–93, [arXiv:1201.5277].
- [8] M. Feshbach, *The integral cohomology rings of the classifying spaces of $O(n)$ and $SO(n)$* , Indiana Univ. Math. J. **32** (1983), no. 4, 511–516.
- [9] D. S. Freed, *Dirac charge quantization and generalized differential cohomology*, in Surv. Differ. Geom. **VII**, 129–194, Int. Press, Somerville, MA, 2000, [arXiv:hep-th/0011220].
- [10] M. B. Green and J. H. Schwarz, *Anomaly cancellation in supersymmetric $D = 10$ gauge theory and superstring theory*, Phys. Lett. **B 149** (1984) 117–122.
- [11] A. Hatcher, *Algebraic Topology*, Cambridge University Press, 2002.

- [12] S.-T. Hu, *Homotopy Theory*, Academic Press, New York, 1959.
- [13] M. Kervaire, *Some nonstable homotopy groups of Lie groups*, Illinois J. Math. **4** (1960), 161–169.
- [14] T. P. Killingback, *Global anomalies, string theory and space-time topology*, Class. Quant. Grav. **5** (1988) 1169–1186.
- [15] A. Kono, *On the integral cohomology of $B\mathrm{Spin}(n)$* , J. Math. Kyoto Univ. **26-3** (1986) 333–337.
- [16] H. B. Lawson, Jr. and M.-L. Michelsohn, *Spin Geometry*, Princeton University Press, Princeton, New Jersey, 1989.
- [17] A. Lichnerowicz, *Spineurs harmoniques*, C. R. Acad. Sci. Paris **257** (1963), 7–9.
- [18] A. T. Lundell, *Concise tables of James numbers and some homotopy of classical Lie groups and associated homogeneous spaces*, Lecture Notes in Math. **1509**, 250–272, Springer, Berlin 1992.
- [19] M. Mimura and H. Toda, *Topology of Lie Groups, I and II*, Amer. Math. Soc., Providence, RI, (1991).
- [20] Th. Nikolaus, C. Sachse, and C. Wockel, *A smooth model for the string group*, Hamburger Beitrage zur Mathematik (2011), no. 405.
- [21] T. Püttmann and A. Rigas, *Presentations of the first homotopy groups of the unitary groups*, Comment. Math. Helv. **78** (2003), 648–662.
- [22] D. Quillen, *The mod 2 cohomology rings of extra special 2-groups and the spinor groups*, Math. Ann., **194** (1971), 197–212.
- [23] C. Redden, *String structures and canonical three-forms*, Pacific J. Math. **249** (2) (2011), 447–484, [arXiv:0912.2086] [math.DG].
- [24] H. Sati, *Twisted topological structures related to M-branes II: Twisted Wu and Wu^c structures* Int. J. Geom. Methods Mod. Phys. **09** (2012), 1250056, [arXiv:1109.4461].
- [25] H. Sati, *Framed M-branes, corners, and topological invariants*, [arXiv:1310.1060] [hep-th].
- [26] H. Sati, *Ninebrane structures*, Int. J. Geom. Methods Mod. Phys. **12** (2015) 1550041, [arXiv:1405.7686] [hep-th].
- [27] H. Sati, U. Schreiber, and J. Stasheff, *Fivebrane structures*, Rev. Math. Phys. **21** (2009), 1197–1240, [arXiv:0805.0564] [math.AT].
- [28] H. Sati, U. Schreiber, and J. Stasheff, *Twisted differential string and fivebrane structures*, Comm. Math. Phys. **315** (2012), no. 1, 169–213, [arXiv:0910.4001] [math.AT].
- [29] H.-B. Shim, *Indefinite string structures*, Ph.D. thesis, University of Pittsburgh, 2013.
- [30] S. Stolz, *A conjecture concerning positive Ricci curvature and the Witten genus*, Math. Ann. **304** (1996), no. 4, 785–800.

- [31] S. Stolz and P. Teichner, *What is an elliptic object?*, in *Topology, geometry and quantum field theory*, 247–343, Cambridge Univ. Press, Cambridge, 2004.
- [32] E. Thomas, *On the cohomology groups of the classifying space for the stable spinor groups*, *Bol. Soc. Mat. Mexicana* (2) **7** (1962) 57–69.
- [33] A. Trautman, *Double covers of pseudo-orthogonal groups*, *Clifford Analysis and Its Applications*, eds. Brace et al., 377–388, Springer, Netherlands, 2001.
- [34] V. V. Varlamov, *Universal coverings of the orthogonal groups*, *Advances in Applied Clifford Algebras* **14**(1) (2004), 81–168.
- [35] K. Waldorf, *String connections and Chern-Simons theory*, *Trans. Amer. Math. Soc.* **365** (2013), 4393–4432, [[arXiv:0906.0117](#)] [[math.DG](#)].
- [36] K. Waldorf, *String geometry vs. spin geometry on loop spaces*, [[arXiv:1403.5656](#)] [[math.DG](#)].
- [37] B.-L. Wang, *Geometric cycles, index theory and twisted K-homology*, *J. Noncommut. Geom.* **2** (2008), no. 4, 497–552.
- [38] G. Whitehead, *Fiber spaces and the Eilenberg homology groups*, *Proc. National Acad. Sciences* **38**, no. 5 (1952), 426–430.
- [39] G. Whitehead, *Elements of homotopy theory*, Springer-Verlag, New York-Berlin, 1978.
- [40] E. Witten, *The index of the dirac operator in loop space*, in *Elliptic Curves and Modular Forms in Algebraic Topology*, *Lecture Notes in Mathematics* **1326** (1988), 161–18.
- [41] J. A. Wolf, *Spaces of constant curvature*, Publish or Perish, Inc. Berkeley, 1977.